



Diversity of bacteria producing pigmented colonies in aerosol, snow and soil samples from remote glacial areas (Antarctica, Alps and Andes)

E. González-Toril, R. Amils, Robert Delmas, J.-R. Petit, J. Komárek, J. Elster

► To cite this version:

E. González-Toril, R. Amils, Robert Delmas, J.-R. Petit, J. Komárek, et al.. Diversity of bacteria producing pigmented colonies in aerosol, snow and soil samples from remote glacial areas (Antarctica, Alps and Andes). Biogeosciences Discussions, 2008, 5 (2), pp.1607-1630. hal-00330295

HAL Id: hal-00330295

<https://hal.science/hal-00330295>

Submitted on 14 Oct 2008

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Biogeosciences Discussions is the access reviewed discussion forum of *Biogeosciences*

Diversity of bacteria producing pigmented colonies in aerosol, snow and soil samples from remote glacial areas (Antarctica, Alps and Andes)

E. González-Toril¹, R. Amils^{1,4}, R. J. Delmas³, J.-R. Petit³, J. Komárek², and J. Elster²

¹Centro de Astrobiología (INTA-CSIC) Ctra. Ajalvir Km. 5, 28850 Torrejón de Ardoz, Spain

²Institute of Botany, Academy of Sciences of the Czech Republic, Trebon and Faculty of Science, University of South Bohemia, Ceske Budejovice, Czech Republic

³Laboratoire de Glaciologie et Géophysique de l'Environnement du CNRS, St Martin d'Heres, France

⁴Centro de Biología Molecular (UAM-CSIC) Cantoblanco, 28049 Madrid, Spain

Received: 12 February 2008 – Accepted: 27 February 2008 – Published: 15 April 2008

Correspondence to: E. González-Toril (glezte@inta.es)

Published by Copernicus Publications on behalf of the European Geosciences Union.

BGD

5, 1607–1630, 2008

**Diversity of
pigmented bacteria in
aerosol, snow and
soil**

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Four different communities and one culture of pigmented microbial assemblages were obtained by incubation in mineral medium of samples collected from high elevation snow in the Alps (Mt. Blanc area) and the Andes (Nevado Illimani summit, Bolivia), from Antarctic aerosol (French station Dumont d'Urville) and a maritime Antarctic soil (King George Island, South Shetlands, Uruguay Station Artigas). Molecular analysis of more than 200 16S rRNA gene sequences showed that all cultured cells belong to the *Bacteria* domain. The phylogenetic comparison with the currently available rDNA database allowed the identification of sequences belonging to *Proteobacteria* (*Alpha*-, *Beta*- and *Gamma*-*proteobacteria*), *Actinobacteria* and *Bacteroidetes* phyla. The Andes snow culture was the richest in bacterial diversity (eight microorganisms identified) and the maritime Antarctic soil the poorest (only one). Snow samples from Col du midi (Alps) and the Andes shared the highest number of identified microorganisms (*Agrobacterium*, *Limnobacter*, *Aquiflexus* and two uncultured *Alphaproteobacteria* clones). These two sampling sites also shared four sequences with the Antarctic aerosol sample (*Limnobacter*, *Pseudonocardia* and an uncultured *Alphaproteobacteria* clone). The only microorganism identified in the maritime Antarctica soil (*Brevundimonas* sp.) was also detected in the Antarctic aerosol. The two snow samples from the Alps only shared one common microorganism. Most of the identified microorganisms have been detected previously in cold environments (*Dietzia kujamenisi*, *Pseudonocardia Antarctica*, *Hydrogenophaga palleronii* and *Brevundimonas* sp.), marine sediments (*Aquiflexus balticus*, *Pseudomonas pseudoalkaligenes*, *Pseudomonas* sp. and one uncultured *Alphaproteobacteria*), and soils and rocks (*Pseudonocardia* sp., *Agrobacterium* sp., *Limnobacter* sp. and two uncultured *Alphaproteobacteria* clones). Air current dispersal is the best model to explain the presence of very specific microorganisms, like those used in this work, in very distant environments. In addition these microorganisms have to be resistant to extreme conditions and able to grow in oligotrophic environments. Considering the habitats in which they have been identified, the presence

BGD

5, 1607–1630, 2008

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of pigments must be related with their ability to resist high doses of radiation.

1 Introduction

Long distance dispersal of biological particles produced by atmospheric circulation, ocean currents, birds, fish, mammals and human vectors, has been known for a long time (Gislén, 1948; Gregory, 1967; Schnell and Vali, 1972; Marshall 1996a, b; Vincent, 2000). Microorganisms have small dimensions (usually less than 5 μm in size) and therefore are easily transported by air masses. Eventually, these wind-born particles are deposited on the ground and/or snow/ice, on very high mountains and polar regions, respectively, in either dry or wet form (Chalmers et al., 1996). It has been proposed that microorganisms living in hot and/or cold terrestrial deserts are especially important for the dispersion of microorganisms due to their special adaptations to extreme and variable conditions (temperature, light intensity, spectral quality, water, desiccation, etc.) (Flechtner, 1999; Van Thielen and Garbary, 1999; Garty, 1999; Elster and Benson, 2004).

Living microorganisms have been collected in the stratosphere (Imshenetsky et al., 1978), and there are several reports supporting the idea that microorganisms can live and reproduce on airborne particles (Dimmick et al., 1979). It has been also shown that microorganisms can actively grow and reproduce at temperatures near or below 0°C, in cloud droplets collected at high altitudes (Sattler et al., 2001).

Both, the viable and dead cells, or their remnants, stored in the ice of continental icecaps and mountain glaciers have the potential to represent a historical record of the recent evolution of microbial life, as well as a record of the Earth's changing climate. In the wake of the recent discovery of the sub-glacial Antarctic Lake Vostok (Priscu et al., 1999; Siegert et al., 2001), and the possibility of recovering water samples containing fossil living microorganisms, there is a growing interest in investigating the transport of living organisms via air over large distances, e.g. cold glacial regions.

In collaboration with the Laboratoire de Glaciologie et Géophysique de

BGD

5, 1607–1630, 2008

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



l'Environnement du CNRS, St Martin d'Heres, France, the composition of micro-autotrophs (cyanobacteria and algae), micro-fungi (hyphae and spores), bacteria (rods, cocci and pigmented bacterial colonies), yeast and plant pollen in remote aerosol, deposited snow, and ice have been recently evaluated (Elster et al., 2007). As a product of this study cultivable pigmented colonies were obtained from alpine snow (Alps and Andes) and aerosol (Antarctic) samples.

It has been suggested (Imshensetsky et al., 1978; Christner et al., 2000) that the presence of highly pigmented bacterial colonies in the mesosphere and in glacial ice was associated with the necessity to protect cells from harmful UV-radiation during atmospheric transport and exposure on the surface of the glaciers. Morphologically similar bacterial specimens have been also observed in soils in the Arctic Svalbard (Rehakova et al., 2008¹). These observations support the idea that this type of microorganisms commonly develop in cold desert ecosystems and are easily transported via aerosol to both, short and long distances.

In this work we report the identification, using molecular ecology methodologies, of the microorganisms able to produce pigmented colonies in enrichment cultures of samples obtained from alpine snow (Alps and Andes) and aerosol (Antarctic). For comparison the culture of pigmented bacteria isolated from maritime Antarctic soil (King George Island, South Shetland) was also analysed.

2 Material and methods

2.1 Site description and sample collection

Aerosol samples originated from a set of filters collected between 1994 and 2000 at the coastal Antarctic Station Dumont d'Urville (66°40' S, 140°01' E). Samples were col-

¹Rehakova, K., Brynychova, K., and Elster, J.: Diversity and abundance of soil algae along the East Brogger deglaciated moraine, Ny-Alesund, Svalbard, Polar Res., submitted, 2008.

BGD

5, 1607–1630, 2008

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



lected on Gelman Zefluor[®] filters (47 mm diameter, 0.5 μ m pore size) by drawing in air at a flow rate of 1.5 m³ h⁻¹. The sampling intervals were of 40 h, which corresponds to a volume between 30 and 60 m³ of air.

In the year 2000, two European alpine snow pits were dug at very high elevation sites in the Mt. Blanc area of the Alps (0.6 m deep) at Col du Midi (elev. 3532 m, 17 May); and the other (2 m deep) at Col du Dome (elev. 4250 m, 31 August). Snow collected from recent deposits contained several visible dust layers. This dust is known to be transported by wind to the Alps from the Sahara desert (Oeschger et al., 1977; Wagenbach, 1989; De Angelis and Gaudichet, 1991). In addition, also in 2000, surface snow blocks were collected at the summit of Nevado Illimani, Bolivia (16°37' S, 67°46' W, Cordillera Real, elevation 6350 m) in the Andes. The Bolivian Andes are surrounded by the Altiplano, a high altitude desert (mean elevation: 3700 m a.s.l., Clapperton, 1993). This region, in particular, contains large salt flats, which are an important source of dust for the regional atmosphere during the dry season (Risacher, 1992). In the austral summer season of 2005, soil samples were collected in close vicinity of the Uruguay Antarctic Station Artigas, King George Island, South Shetlands.

All samples were collected using pre-cleaned glass vials (aerosol) and sealed plastic bags (snow and soil), respectively, transported frozen to the laboratory in Grenoble and/or in Tebo and stored frozen until further analysis (for more details see Elster et al., 2007).

2.2 Sample preparation and cultivation

To minimise possible contaminations, all post-sampling manipulations were done in a UV-sterilised laminar box hood, using sterile glass vials. To retrieve samples from the snow pits an upper layer of about 2 cm was sliced with a sharp knife. Snow samples were dried off at a temperature of -40°C in a special sterile lyophilisation device (Lyovac GT2, Leybold-Heraeus, Germany). After this operation, glass vessels containing solid deposits were rinsed with 10–15 ml of re-distilled water.

BGD

5, 1607–1630, 2008

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The aerosol filters were cut into quarters and one quarter was used for enrichment cultures. Each snow and aerosol sample was cultured in a sterile glass bottle (25 ml in volume). The bottom of each glass bottle was filled to a depth of one centimeter with glass beads (diameter 3 mm). BG-11 culture media (Bischoff and Bold, 1963) was used for all experiments. Each glass bottle was filled with about 10 ml of sterile medium (for more details see Elster et al., 2007). For soil samples analysis, suspensions of 10 g soil + 90 ml sterile water were homogenized using ultrasonication for 4 min and spread on Petri dishes containing mineral agar (0.1 ml of soil suspension per Petri dish with 1.5% BG11 agar in four replicates) (Elster et al., 1999). Glass bottles and Petri dishes were cultivated in an illuminated ($\sim 100 \text{ W cm}^2$) refrigerator (temperature 5–8°C) with a light regime of 18 h of light, 2 h of UV-B radiation (germicide lamp) and 4 h of darkness. The germicide lamp was used to sterilize the culture growth area (UV-B light did not penetrate through the glass bottles). Experimental bottles were shaken every 2–3 days. After 1 and 2 months of cultivation, the contents of bottles and Petri dishes were analyzed under the light microscope (Olympus BX 60). Selected samples with pigmented microbial colonies were analyzed by fluorescence after staining with the DAPI fluorochrome (EFM Olympus BX 60) (Zachleder and Cepák, 1987) and transmission electron microscopy (Jeol equipment JEM 1010).

Five pigmented microbial colonies were selected for further analysis (Fig. 1):

1. aerosol sample collected on 7 July 1999, in the coastal Antarctic Station Dumont d'Urville (66°40' S, 140°01' E).
2. Col du Midi (Alps) snow samples from a depth between 10 to 20 cm (elevation. 3532m, collected on 17 May 2000).
3. Col du Dome, Vallot (Alps) snow samples from surface to a depth of 10 cm (elevation. 4250 m, collected on 31 August 2000).
4. Nevado Illimani (Andes) surface snow blocks (16°37' S, 67°46' W, Cordillera Real, elevation 6350 m) collected in 2000.

BGD

5, 1607–1630, 2008

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5. Soil sample from the Uruguay Antarctic Station Artigas, King George Island, South Shetland Islands (62°10' S, 58°30' W), collected in the austral summer of 2005.

All cultures were transported to the Centro de Astrobiología (CAB) in Madrid, where microorganisms were identified by comparison of 16S rRNA gene sequences.

2.3 DNA extraction

One ml of each enrichment culture was used for DNA extraction. Fast DNA Spin kit for soil (Q-Bio Gene Inc., CA, USA) was used according to the manufacturer's instructions. To disrupt the cells, the mixture of ceramic and silica beads provided in the kit and three pulses of 40 s at speed 5.5 of the FastPrep bead-beating instrument (Bio 101) were applied. After the extraction, DNA was purified by passage through a GeneClean Turbo column (Q-Bio Gene Inc., CA, USA) and quantified by ethidium bromide-UV detection on an agarose gel (González-Toril et al., 2006).

2.4 16S ribosomal RNA clone library construction

PCR amplification of 16S rRNA gene fragments between *E. coli* positions 8 and 1507 for *Bacteria* domain (Lane, 1991), between *E. coli* position 25 and 1492 for *Archaea* domain (Achenbach, and Woese, 1995), and between *E. coli* position 359 and 805 for *Cyanobacteria* (Nübel et al., 1997) were performed. These genes were amplified by PCR in mixtures containing 20-30 ng of DNA per 50 μ l reaction volume, 1 \times PCR buffer (Promega Biotech Iberica, Spain), 2.5 μ M of each of the deoxynucleotides (Amersham Biosciences, UK), 2.5 mM MgCl₂, 1 mg mL⁻¹ bovine serum albumin (BSA), 500 mM of each forward and reverse primers and 0.025 U/ μ l of Taq DNA polymerase (Promega Biotech Iberica, Spain) (Table 1). PCR conditions were as follows: initial denaturation at 94°C for 5 min, followed by 30 cycles of denaturation at 94°C for 40 s, annealing at 52°C for 1 min for the *Bacteria* domain, 56°C for the *Archaea* domain, and 60°C for the *Cyanobacteria*. Large 16S rRNA gene fragments (>1400 bp) were purified by

BGD

5, 1607–1630, 2008

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



GeneClean Turbo Column (Q-Bio Gene Inc. CA, USA) and cloned using the Topo Ta Cloning Kit (Invitrogen. CA, USA). Cloned inserts were amplified using PCR conditions described above and they were directly sequenced with a Big-Dye sequencing kit (Applied Biosystem) following the manufacturer's instructions (González-Toril et al., 2006).

2.5 Clone library analysis

Sequences were analyzed using BLAST at the NCBI database (<http://ncbi.nlm.nih.gov/BLAST>) and added together with the most important BLAST hits, to an alignment of 50 000 homologous bacterial 16S rRNA primary structures by using the aligning tool of the ARB software package (<http://www.arb-home.de>) (Ludwig et al., 2004). The rRNA alignments were corrected manually and alignment uncertainties were omitted in the phylogenetic analysis. Phylogenetic trees were generated using parsimony, neighbour-joining, and maximum-likelihood analyses with a subset of 200 nearly full-length sequences (>1400 bp). Filters, which excluded highly variable positions, were used. In all cases, general tree topology and clusters were stable. A consensus tree was generated (González-Toril et al., 2006). Sequences obtained in this study have been deposited in the EMBL sequence database under accession numbers from EU429484 to EU429508.

3 Results and discussion

Following the protocols for enrichment cultures of photoautotrophic microorganisms present in snow samples from the Alps and the Andes and in an aerosol sample from Antarctica, different pigmented non-photosynthetic prokaryotes were able to grow in the extremely poor nutritional conditions of the selected media (Fig. 1) (Elster et al., 2007). Due to the characteristics of the cultures and the idiosyncrasy of the samples it was considered of interest to identify the microorganisms present in the different cultures, to

BGD

5, 1607–1630, 2008

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

evaluate the diversity corresponding to this very specific type of microorganisms, and eventually to compare the results obtained in geographically dispersed sampling sites. A commercial soil DNA extraction kit was used to extract DNA from the enrichment cultures from particulate matter present in high elevation snow in the Alps and the Andes, from an Antarctic aerosol and a maritime Antarctic soil. DNA was extracted from all samples. No phototrophic *Bacteria* (cyanobacteria) were amplified using specific primers in any of the DNA extracted from the five samples. These results agree with the microscopy analysis of the cultures. Also no *Archaea* were detected by amplification.

PCR amplification of 16S rRNA genes using universal bacterial primers, followed by cloning and sequencing produced around 200 sequences which were used to identify the different bacteria present in the enrichment cultures by comparison with the NCBI database. The retrieved sequences were added to a database of over 50 000 prokaryotic 16S rRNA gene sequences using the aligning tool of the ARB software package (<http://www.arb-home.de>) (Ludwig et al., 2004). Phylogenetic trees were generated using parsimony, neighbour joining, and maximum-likelihood analyses with a subset of 200 nearly full-length sequences (>1400 bp). Filters excluding highly variable positions were used. In all cases general tree topology and clusters were stable. Thus, consensus trees were generated (Figs. 2–6).

After the analysis of the generated sequences representatives of three bacterial phyla: *Proteobacteria* (*Alpha*, *Beta* and *Gammaproteobacteria*), *Actinobacteria* and *Bacteroidetes*, were identified (Fig. 2). Seven close relative microorganisms have been identified for the snow sample from Col du Midi (Alps). Three sequences corresponded to the *Alphaproteobacteria*: one as a possible member of the genus *Agrobacterium* (closest relative: clone IrT-J614-14, retrieved from an environmental sample of a uranium mine (Selenska-Pobell, 2002)) (Fig. 3); the other two, close to clones B3NR69D12 and AP-12, detected in a underground cave (Northup et al., 2003) and in a marine estuary (NCBI accession number AY145551, unpublished) respectively. Two sequences corresponded to the *Betaproteobacteria*: one a possible member of the species *Hydrogenophaga palleronii* (Fig. 4), identified by FISH in a lake snow ag-

**Diversity of
pigmented bacteria in
aerosol, snow and
soil**

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

gregates (Schweitzer et al., 2001), and the other close to a clone of the genus *Limnobacter*, D-15, isolated from underground mineral water (Loy et al., 2005). One sequence corresponding to the *Gammaproteobacteria* exhibited a high homology with *Pseudomonas pseudoalkaligenes*, which was isolated from a marine sediment (NCBI accession number AF286035, unpublished). One sequence corresponded to the phylum *Bacteroidetes*, with high homology with *Aquiflexus balticus* (Fig. 5), also retrieved from a marine sediment (Brettar et al., 2004).

The other sample from the Alpes (Col du Dome) exhibited a much lower diversity than the one obtained from Col du Midi. In this sample only three sequences were retrieved. Two corresponded to uncultivated *Alphaproteobacteria* clones, AP-12 and O15B-H01, the first detected in a marine estuary (NCBI accession number AY145551, unpublished) and the other from ground water of a uranium mine (NCBI accession number AY662032, unpublished). The other sequence corresponded to the class *Actinobacteria*, with a high level of homology with the species *Dietzia kujamensis*, which was isolated from the Himalaya (Mayilraj et al., 2006).

Eight close relative microorganisms have been identified for the snow sample from the Andes (Nevado Illimani). Three sequences corresponded to the *Alphaproteobacteria* class: two related to the *Agrobacterium* (clone IrT-J614-14) (Fig. 3) and the clone B3NR69D12, both identified in Col du Midi; and one related to the uncultured EV818CFSSAHH29 clone, which has been identified in subsurface water of the Kalahari Shield in South Africa (NCBI accession number DQ336984, unpublished). Two sequences corresponded to the *Betaproteobacterai* class: *Hydrogenophaga palleronii* (Fig. 4) and *Limnobacter* clone D-15, both also identified in Col du Midi. One sequence exhibiting high homology with *Aquiflexus balticus* (Fig. 5) of the phylum *Bacteroidetes*, already described in Col du Midi, has been retrieved from the Andes enrichment culture. The last two sequences corresponded to the *Actinobacteria* class: one with high homology to *Microbacterium thalassium* (Richert, 2007) and the other with *Pseudonocardia* sp. (Fig. 6) which has been isolated from marine sediments (NCBI accession number AY974793, unpublished).

BGD

5, 1607–1630, 2008

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seven sequences retrieved from the aerosol sample from Antarctica were identified by phylogenetic analysis. Three sequences corresponded to the *Alphaproteobacteria* class: one related with clone B3NR69D12, which has been also identified in the Andes and the Alps (Col du Midi); another related with clone AP-12 which has been also identified in Col du Midi; and a third one corresponding to the *Brevundimonas vesicularis*, which was isolated from an ice core of Greenland (Sheridan et al., 2003). One sequence corresponded to the *Betaproteobacteria* and had a high level of homology with the uncultured clone of *Limnobacter* D-15, already identified in the enrichment cultures of snow samples from the Andes and the Alps (Col du Midi). Three sequences corresponded to the *Actinobacteria* class: one related with *Pseudonocardia antarctica* (Fig. 3), a psychrophilic microorganisms isolated from McMurdo Valley in the Antarctica (Prabakar et al., 2004); a second one corresponding to a member of the *Pseudonocardia* genus (Fig. 6) isolated from marine sediments and also identified in the Andes; and a third one similar to *Brachybacterium conglomeratum*, which has been isolated from a spacecraft assembly facility (NCBI accession number AY145551, unpublished).

The sample from maritime Antarctica soil was a pure culture because only one sequence was retrieved with a high level of 16S rRNA gene sequence homology with a member of the *Brevundimonas* sp. identified previously in permafrost samples and close to the *Alphaproteobacteria* species of *Brevundimonas* identified in the Antarctic aerosol.

All the identified microorganisms in the enrichment cultures belong to the bacterial domain. Snow samples from the Alps (Col du Midi) and the Andes (Illimani), and the Antarctic aerosol sample exhibited a similar level of bacterial diversity in the corresponding enrichment cultures (Table 2). The maritime Antarctic soil (King George Island, Uruguay Antarctic Station Artigas) enrichment culture contained only one sequence (Table 2). This is probably due to the protocol used for its enrichment. Only one type of bacteria was able to grow on mineral agar plate. On the contrary, in the case of snow and aerosol samples growth in liquid mineral media was always observed. The

**Diversity of
pigmented bacteria in
aerosol, snow and
soil**

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

distribution of detected bacteria was related with cold environments (21.4%), marine environments (35.7%) and soils/subsurface (35.7%) (Table 2). No single cosmopolitan bacterium was found in all the analyzed sites, although three geographically distant sampling sites (Alps, Illimani and Antarctic aerosol) shared three microorganisms (*Lim-*
nobacter clone D-15, and two uncultured *Alphaproteobacteria*, clones B3NR69D12 and AP-12). Different bacteria were found in two locations: *Agrobacterium* sp. (Col du Midi and Illimani), *Brevundimonas* sp. (Antarctic aerosol and maritime Antarctic soil), *Hydrogenophaga palleroni* (Col du Midi and Illimani), *Aquiflexus balticus* (Col du Midi and Illimani) and *Pseudonocardia* sp. (Illimani and Antarctic aerosol) (Table 2). Six out
of fourteen identified genus or species are found only in a single location (Table 2). Interestingly enough the diversity found in very close locations in the Alps (Col du Dome and Col du Midi) are much more different (only one common sequence out of ten) than the diversity found in locations in different hemispheres, altitudes or matrix (snow versus aerosol) (Table 2). Concerning the pigments giving color to the microor-
ganisms, several bacteria related with those present in the enrichment cultures have been described as having this peculiar phenotypic property, like *Dietzia*, *Aquiflexum*, *Microbacterium* and *Pseudonocardia*. In addition, members of the *Actinobacteria* class are well known by their ability to synthesize pigments for radiation protection purposes.

Concerning the different models for long distance microbial dissemination (atmo-
spheric circulation, ocean currents, birds, fish, mammals and human vectors), if we consider the habitats in which related microorganisms have been detected, we can rule out ocean currents and animal vectors, leaving atmospheric circulation as the most plausible mean of dissemination, specially considering the microbial diversity detected in the Antarctic aerosol and shared with other distant locations (71.4%), which underlines the importance of microbes attached to dust particles for its disseminations.

Of course our data can not rule out the possible model of “everything is everywhere”, although considering the experimental constraints introduced by the use of extremely oligotrophic selective media, we think that atmospheric dispersion can explain adequately the observed results. One of the problems related with this type of experiments

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

is to rule out the possibility of contamination. Considering the location of the selected sampling sites and the obtained results we think that this possibility is improbable. Although the difference observed between the samples obtained in the same campaign in two close sites in the Alps is not easy to explain, it is a very useful internal control of lack of contamination on the sampling manipulation. We strongly believe that the common microbial patterns of peculiar microorganisms observed at distant locations reflects truly airborne microbial dissemination, so we will like to propose this protocol for further testing this model of microbial dispersion.

Acknowledgements. The work has been supported by grants from the Ministerio de Educación y Ciencia from Spain (grant CGL2006/02534/BOS), the Ministry of Education of the Czech Republic (Kontakt – ME 934, ME 945 and project GA CR 206/05/0253) and French-Czech cooperative research programme Barrande No. 99054. We are very grateful to J. Snokhousova for technical assistance.

References

- Achenbach, L. and Woese, C.: 16S and 23S rRNA-like primers, in: Archaea. A laboratory manual, edited by: Sower, R. and Schreier, H. J., Cold Spring Harbor Laboratory Press, NY, 521–523, 2003.
- Bischoff, H. W. and Bold, H. C.: Some algae from Enchanted Rock and related algal species, in Phycological studies, University of Texas, IV, Austin, 68, 1963.
- Brettar, I., Christen R., and Höfle, M. G.: *Aquiflexum balticum* gen. nov., sp. nov., a novel marine bacterium of the *Cytophaga-Flavobacterium-Bacteroides* group isolated from surface water of the central Baltic Sea, Int. J. Syst. Evol. Microbiol., 54(6), 2335–2341, 2004.
- Brosius, J., Dull, T. J, Sleeter, D. D., and Noller, H. F.: Gene organization and primary structure of a ribosomal RNA operon from *Escherichia coli*, J. Mol. Biol., 148, 107–127, 1981.
- Clapperton, C. C.: Quaternary Geology and Geomorphology of South America, in: Elsevier Science Publ., Amsterdam, 780, 1993.
- Chalmers, M. O., Harper, M. A., and Marshall, W. A.: An Illustrated Catalogue of Airborne Microbiota from the Maritime Antarctic, British Antarctic Survey, Cambridge, 175, 1996.

BGD

5, 1607–1630, 2008

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Christner, B. C., Mosley-Thompson, E., Thompson, L. G., Zagorodnoc, V., Sandman, K., and Reeve, J. N.: Recovery and identification of viable bacteria immured in glacial ice, *Icarus*, 144, 479–485, 2000.

De Angelis, M. and Gaudichet, A.: Saharan dust deposition over Mont Blanc (French Alps) during the last 30 years, *Tellus* 43B, 61–75, 1991.

Dimmick, R. L., Wolochow, M. A., and Chatigny, A. A.: Evidence for more than one division of bacteria within airborne particles. *Appl. Environ. Microbiol.*, 38, 642–643, 1979.

Elster, J., Lukešová, A., Svoboda, J., Kopecký, J., and Kanda, H.: Diversity and abundance of soil algae in the polar desert, Sverdrup pass, central Ellesmere Island, *Polar Rec.*, 35, 231–254, 1999.

Elster, J. and Benson, E. E.: Life in the Polar Terrestrial Environment: A Focus on Algae, in: *Life In The Frozen State*, edited by: Fuller, B., Lane, N., and Benson, E. E., Taylor and Francis, London, 111–149, 2004.

Elster, J., Delmas, R. J., Petit, J.-R., Řeháková, K.: Composition of microbial communities in aerosol, snow and ice samples from remote glaciated areas (Antarctica, Alps, Andes), *Biogeosciences Discuss.*, 4, 1–35, 2007,

<http://www.biogeosciences-discuss.net/4/1/2007/>.

Flechlner, V. R.: Enigmatic desert soil algae. Soil algal flora of the Western U.S.A. and Baja California, Mexico, in: *Enigmatic Microorganisms and Life in Extreme Environments*, edited by: Seckbach, J., Kluwer Academic Publishers, 231–241, 1999.

Garty, J.: Lithobionts in the Eastern Mediterranean, in: *Enigmatic Microorganisms and Life in Extreme Environments*, edited by: Seckbach, J., Kluwer Academic Publishers, 255–276, 1999.

Gislén, T.: Aerial plankton and its conditions of life, *Biological Reviews*, 23, 109–126, 1948.

González-Toril, E., Gómez, F., Malki, M., and Amils, R.: The isolation and study of acidophilic microorganisms, in: *Extremophiles – Methods in Microbiology*, edited by: Rainey, F. A. and Oren, A., Elsevier/Academic Press, Amsterdam, The Netherlands, 35, 471–510, 2006.

Gregory, P. H.: Atmospheric microbial cloud systems, *Science Progress*, Oxford, 55, 613–628, 1967.

Imshenetsky, A. A., Lysenko, S. V., and Kazakov, G. A.: Upper boundary to the biosphere, *Appl. Environ. Microbiol.*, 35, 1–5, 1978.

Lane, D. J.: 16S/23S rRNA sequencing, in: *Nucleic acid techniques in bacterial systematics*, edited by: Stackebrandt, E. and Goodfellow, M., John Wiley & Sons Ltd., Chichester, UK,

BGD

5, 1607–1630, 2008

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1991.

Loy, A., Beisker, W., and Meier, H.: Diversity of bacteria growing in natural mineral water after bottling, *Appl. Environ. Microbiol.*, 71(7), 3624–3632, 2005.

Ludwig, W., Strunk, O., Westram, R., Richter, L., Meier, H., Yadhukumar, Buchner, A., Lai, T., Steppi, S., Jobb, G., Förster, W., Brettske, I., Gerber, S., Ginhart, A. W., Gross, O., Grumann, S., Hermann, S., Jost, R., König, A., Liss, T., Lüßmann, R., May, M., Nonhoff, B., Reichel, B., Strehlow, R., Stamatakis, A., Stuckmann, N., Vilbig, A., Lenke, M., Ludwig, T., Bode, A., and Schleifer, K. H.: ARB: a software environment for sequence data, *Nucleic Acids Res.*, 32(4), 1363–1371, 2004.

10 Marshall, W. A.: Aerial dispersal of lichens soredia in the maritime Antarctic, *New Phytologist*, 134, 523–530, 1996a.

Marshall, W. A.: Biological particles over Antarctica, *Nature*, 383, 680 pp., 1996b.

Mayilraj, S., Suresh, K., Kroppenstedt, R. M., and Saini, H. S.: *Dietzia kunjamensis* sp. nov., isolated from the Indian Himalayas, *Int. J. Syst. Evol. Microbiol.*, 56(7), 1667–1671, 2006.

15 Northup, D. E., Barns, S. M., Yu, L. E., Spilde, M. N., Schelble, R. T., Dano, K. E., Crossey, L. J., Connolly, C. A., Boston, P. J., Natvig, D. O., and Dahm, C. N.: Diverse microbial communities inhabiting ferromanganese deposits in Lechuguilla and Spider Caves, *Environ. Microb.*, 5(11), 1071–1086, 2003.

20 Nübel, U., Garcia-Pichel, F., and Muyzer, G.: PCR primers to amplify 16S rRNA genes from cyanobacteria, *Appl. Environ. Microb.*, 63, 3327–3332, 1997.

Oeschger, H., Schotterer, U., Stauffer, B., Haeberli, W., and Röthlisberger, H.: First results from Alpine core drilling projects, *Ztschr. Gletscherk. Glazialgeolog.*, 13(1/2), 193–208, 1977.

25 Prabahar, V., Dube, S., Reddy, G. S., and Shivaji, S.: *Pseudonocardia antarctica* sp. nov. an *Actinomycetes* from McMurdo Dry Valleys, Antarctica, *Syst. Appl. Microbiol.*, 27, 66–71, 2004.

Priscu, J. C., Adams, E. E., Lyons, W. B., Voytek, M. A., Mogk, D. W., Brown, R. L., McKay, C. P., Takacz, C. D., Welch, K. A., Wolf, F. C., Kirshtein, J. D., and Acci, R.: Geomicrobiology of subglacial ice above lake Vostok, Antarctica, *Science*, 286, 2141–2144, 1999.

30 Richert, K., Brambilla, E., and Stackebrandt, E.: The phylogenetic significance of peptidoglycan types: Molecular analysis of the genera *Microbacterium* and *Aureobacterium* based upon sequence comparison of *gyrB*, *rpoB*, *recA* and *ppk* and 16SrRNA genes, *Syst. Appl. Microb.*, 30(2), 102–108, 2007.

Risacher, F.: Géochimie des lacs salés et croûtes de sel de l'Altiplano Bolivien., *Sci. Geol. Bull.*,

BGD

5, 1607–1630, 2008

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- 45, 135–214, 1992.
- Sattler, B., Puxbaum, H., and Psenner, R.: Bacterial growth in supercooled cloud droplets, *Geophys. Res. Lett.*, 28(2), 239–242, 2001.
- Schnell, R. C. and Vali, G.: Atmospheric ice nuclei from decomposing vegetation, *Nature*, 236, 163–165, 1972.
- Schweitzer, B., Huber, I., Amann, R., Luwig, W., and Simon, M.: α - and β -*Proteobacteria* control the consumption and release of amino acids on lake snow aggregates, *Appl. Environ. Microbiol.*, 67(2), 632–645, 2001.
- Selenska-Pobell, S.: Diversity and activity of bacteria in uranium mining waste piles, in: *Radioactivity in the environment, Interactions of microorganisms with radionuclides*, edited by: Keith-Roach, M. J. and Lievens, F. R., Elsevier, 22, 225–254, 2002.
- Sheridan, P. P., Miteva, V. I., and Brenchley, J. E.: Phylogenetic analysis of anaerobic psychrophilic enrichment cultures obtained from a Greenland glacier ice core, *Appl. Environ. Microbiol.*, 69(4), 2153–2160, 2003.
- Siebert, M. J., Ellis-Evans, J. C., Tranter, M., Mayer, C., Petit, J. R., Salamatin, A., and Prisco, J. C.: Physical, chemical and biological processes in Lake Vostok and other Antarctic subglacial lakes, *Nature*, 414, 603–609, 2001.
- Van Thienen, N. and Garbary, D. J.: Life in the rocks – endolithic algae, in: *Enigmatic Microorganisms and Life in Extreme Environments*, Seckbach, J., Kluwer Academic Publishers, 243–253, 1999.
- Vincent, W. F.: Evolutionary origins of Antarctic microbiota: invasion, selection and endemism, *Antarct. Sci.*, 12(3), 374–385, 2000.
- Wagenbach, D.: Environmental records in Alpine glaciers, in: *The environmental records in glaciers and ice sheets*, edited by: Oeschger, J. H. and Langway, C. C., John Wiley & Sons Limited, 69–83, 1989.
- Zachleder, V. and Cepák, V.: Visualisation of DNA containing structures by fluorochrome DAPI in those algal cells that are not freely permeable to the dye, *Arch. Hydrobiol./Algolog. Stud.*, 47, 157–168, 1987.

BGD

5, 1607–1630, 2008

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.

Table 1. Primers used for PCR amplification.

Primer ^a	Target site ^b	Sequence (5' to 3')	Specificity	Reference
8F	8–23	AGA GTT TGA TCM TGG C	<i>Bacteria</i> Domain	Lane (1991)
25F	9–25	CYG GTT GAT CCT GCC RG	<i>Archaea</i> Domain	Achenbach and Woese (1995)
1492R	1492–1513	TAC GGY TAC CTT GTT ACG ACT T	Universal	Achenbach and Woese (1995)
Cya359F	359–378	GGG GAA TYT TCC GCA ATG GG	<i>Cyanobacteria</i> and chloroplast	Nübel et al. (1997)
Cya106F	106–127	CGG ACG GGT GAG TAA CGC GTG A	<i>Cyanobacteria</i> and chloroplast	Nübel et al. (1997)
Cya781R(a) ^c	781–805	GAC TAC TGG GGT ATC TAA TCC CAT T	<i>Cyanobacteria</i> and chloroplast	Nübel et al. (1997)
Cya781R(b) ^c	781–805	GAC TAC AGG GGT ATC TAA TCC CTT T	<i>Cyanobacteria</i> and chloroplast	Nübel et al. (1997)

^a F (forward) and R (reverse) indicate the orientations of the primers in relation to the rRNA.

^b Positions are given according to the *E. coli* numbering of Brosius et al. (1981).

^c Reverse primer CYA781R is an equimolar mixture of CYA781R(a) and CYA781R(b).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Closet phylogenetic relatives. Microorganisms and sequences closely related with clones obtained in this studies and some of their characteristics.

Col du Dome (Alps)	Col du Midi (Alps)	Illimani (Andes)	Artigas (Antarctic)	aerosol (Antarctic)	Interesting characteristics	NCBI ^a
Clone AP-12 Near <i>Bradyrhizobium</i> <i>Alphaproteobacteria</i>	Clone AP-12 Near <i>Bradyrhizobium</i> <i>Alphaproteobacteria</i>	—	—	Clone AP-12 Near <i>Bradyrhizobium</i> <i>Alphaproteobacteria</i>	Uncultured bacterium from Weser Estuary	AY145551
—	Clone B3NR69D12 Near <i>Altipia massiliensis</i> <i>Alphaproteobacteria</i>	Clone B3NR69D12 Near <i>Altipia massiliensis</i> <i>Alphaproteobacteria</i>	—	—	Un cultured bacterium from an inhabiting ferromanganese deposits in Lechugilla and Spider Caves	AY186080
Clone O15B-H01 Near <i>Sinella granulii</i> <i>Alphaproteobacteria</i>	—	—	—	—	Uncultured bacterium from a groundwater contaminated with high levels of nitric acid-bearing uranium water	AY662032
—	Clone IrT-J614-14 Near <i>Agrobacterium</i> <i>Alphaproteobacteria</i>	Clone IrT-J614-14 Near <i>Agrobacterium</i> <i>Alphaproteobacteria</i>	—	—	Uncultured bacterium from a uranium mining waste piles	AJ295675
—	—	Clone EV818CFSSH29 Near <i>Brevundimonas</i> <i>Alphaproteobacteria</i>	<i>Brevundimonas</i> sp. <i>Alphaproteobacteria</i>	<i>Brevundimonas vesicularis</i> <i>Alphaproteobacteria</i>	— Uncultured bacterium from subsurface water of Kalarhari Shield, South Africa — Bacteria isolated from permafrost. — Anaerobic psychrophilic enrichment cultures obtained from a Greenland glacier ice core — Bacteria detected in lake snow aggregates by FISH	DQ336984 DQ177489 AY169433
—	Clone D-15 Near <i>Limnobacter</i> <i>Betaproteobacteria</i>	Clone D-15 Near <i>Limnobacter</i> <i>Betaproteobacteria</i>	—	Clone D-15 Near <i>Limnobacter</i> <i>Betaproteobacteria</i>	Uncultured bacterium from natural mineral water	AF522999
—	<i>Hydrogenophaga palleronii</i> <i>Betaproteobacteria</i>	<i>Hydrogenophaga palleronii</i> <i>Betaproteobacteria</i>	—	—	Bacteria detected in lake snow aggregates by FISH	AF078769
—	Clone LCP-79 <i>Pseudomonas</i> <i>Gammaproteobacteria</i>	—	—	—	Uncultured bacterium from marine sediments	AF286035
<i>Dietzia</i> spp. <i>Actinobacteria</i>	—	—	—	—	Bacteria isolated in cold environments (Himalaya and arctic Ocean)	DQ060378 AY972480 AB159036
—	—	<i>Pseudonocardia</i> sp. <i>Actinobacteria</i>	—	<i>Pseudonocardia antarctica</i> <i>Actinobacteria</i>	— Bacteria isolated from marine sediments (depths of 500 m). — Bacteria isolated from McMurdo Dry Valleys, Antarctica. Filamentous and produce brown colour substrate mycelia and aerial mycelia Vich form a white conglomerate.	AY974793 DQ448725 AJ576010
—	—	<i>Microbacterium thalassium</i> <i>Actinobacteria</i>	—	—	In young cultures, cells are small irregular rods. In old cultures, rods become shorter or spherical elements. Colonies are yellowish white, yellow or orange.	AM181507
—	—	—	—	<i>Brachybacterium</i> sp. <i>Actinobacteria</i>	Bacteria isolated from spacecraft assembly facilities	AY167842
—	<i>Aquiflexum balticum</i> <i>Bacteroidetes</i>	<i>Aquiflexum balticum</i> <i>Bacteroidetes</i>	—	—	Bacterium isolated from surface water of the Central Baltic Sea (depth of 5 m). Red and transparent colonies when young, but turn opaque with ongoing incubation. Cells contain carotenoids.	AJ744861

^a Accession Number in NCBI of the closest phylogenetic relatives

BGD

5, 1607–1630, 2008

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Diversity of
pigmented bacteria in
aerosol, snow and
soil**

E. González-Toril et al.



© Elster et al., 2007

Fig. 1. Display of pigmented microorganisms enriched from snow and aerosol samples.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.




Fig. 2. Prokaryotic phylogenetic tree showing in red the phyla in which pigmented microorganisms have been identified.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.




Fig. 3. Phylogenetic tree of *Agrobacterium* and related bacteria. Phylogenetic trees were generated using parsimony, neighbor-joining and maximum likelihood analysis with different sets of filters. All generated phylogenetic trees resulted in stable branching. A consensus tree was generated using all calculated trees. Cloned 16S rRNA sequences from enrichment cultures are indicated in bold. Sequences retrieved from Nevado Illimani (Andes) are designated by the word “Illimani” followed by the number of the sequence. Sequences from Col du Midi (Alps) are designated by “Coldumidi” followed by the number of the sequence. Bar represents 10% estimated phylogenetic divergence.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.




Fig. 4. Phylogenetic tree of *Hydrogenophaga* and related bacteria. Phylogenetic trees were generated using parsimony, neighbor-joining and maximum likelihood analysis with different sets of filters. All generated phylogenetic trees resulted in stable branching. A consensus tree was generated using all calculated trees. Cloned 16S rRNA sequences from enrichment cultures are indicated in bold. Sequences retrieved from Nevado Illimani (Andes) are designated by the word “Illimani” followed by the number of the sequence. Sequences from Col du Midi (Alps) are designated by “Coldumidi” followed by the number of the sequence. Bar represents 10% estimated phylogenetic divergence.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.




Fig. 5. Phylogenetic tree of *Aquiflexus* and related bacteria. Phylogenetic trees were generated using parsimony, neighbor-joining and maximum likelihood analysis with different sets of filters. All generated phylogenetic trees resulted in stable branching. A consensus tree was generated using all calculated trees. Cloned 16S rRNA sequences from enrichment cultures are indicated in bold. Sequences retrieved from Nevado Illimani (Andes) are designated by the word “Illimani” followed by the number of the sequence. Sequences from Col du Midi (Alps) are designated by “Coldumidi” followed by the number of the sequence. Bar represents 10% estimated phylogenetic divergence.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Diversity of pigmented bacteria in aerosol, snow and soil

E. González-Toril et al.




Fig. 6. Phylogenetic tree of *Pseudonocardia* and related bacteria. Phylogenetic trees were generated using parsimony, neighbor-joining and maximum likelihood analysis with different sets of filters. All generated phylogenetic trees resulted in stable branching. A consensus tree was generated using all calculated trees. Cloned 16S rRNA sequences from enrichment cultures are indicated in bold. Sequences retrieved from Nevado Illimani (Andes) are designated by the word “Illimani” followed by the number of the sequence. Sequences from aerosol sample recollected in Antarctica are designated by “Aeroplankton” followed by the number of the sequence. Bar represents 10% estimated phylogenetic divergence. Additional figures can be viewed in the supplementary material (<http://www.biogeosciences-discuss.net/5/1607/2008/bgd-5-1607-2008-supplement.pdf>).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion